

Reclassifications in *Halosarpheia* and related genera with unfurling ascospore appendages

by

K.L. Pang*

School of Biological Sciences, University of Portsmouth, King Henry Building
King Henry I Street, Portsmouth PO1 2DY, UK

and

E.B.G. Jones

National Center for Genetic Engineering and Biotechnology
BIOTEC Central Research Unit
113 Paholyothin Road, Klong 1, Klong Luang, Pathumthani 12120, Thailand

Pang, K.L. & E.B.G. Jones (2004): Reclassifications in *Halosarpheia* and related genera with unfurling ascospore appendages. - Nova Hedwigia 78: 269-271.

That *Halosarpheia* as presently conceived is polyphyletic has been sufficiently demonstrated in recent years (Kong et al. 2000, Abdel-Wahab et al. 2001, Anderson et al. 2001) and a generic reclassification was deemed necessary. In three papers concerned with species of *Halosarpheia* and related genera with polar unfurling ascospore appendages (Campbell et al. 2003, Pang et al. 2003a, b), a subdivision of the genus was proposed. The acceptance and publication dates of these papers overlap and therefore nomenclatural priority for the new names proposed in these papers must be established.

Saagaromyces K.L. Pang & E.B.G. Jones 2003a
≡ *Littispora* J. Campb., J.L. Anderson & Shearer 2003

Saagaromyces abonnis (Kohlm.) K.L. Pang & E.B.G. Jones 2003a
Basionym: *Halosarpheia abonnis* Kohlm. 1984
≡ *Littispora abonnis* (Kohlm.) J. Campb., J.L. Anderson & Shearer 2003

*Corresponding author. E-mail: <ka-lai.pang@port.ac.uk>